

# The Friedman Universe with the stochastic cosmological constant

---

**R. Manka**

*University of Silesia, Institute of Physics, Katowice, Poland*  
*E-mail:manka@us.edu.pl*

**D. Panchyrz**

*University of Silesia, Institute of Physics, Katowice, Poland*

**ABSTRACT:** The Friedman Universe with the remnant stochastic scalar field in the hybrid inflation model is examined. It is shown that the small effective cosmological constant  $\delta\Omega_\Lambda$  appears which increases the cosmological expansion.

**KEYWORDS:** Classical Theories of Gravity, Stochastic Processes.

---

## Contents

|                                   |          |
|-----------------------------------|----------|
| <b>1. Introduction</b>            | <b>1</b> |
| <b>2. The classical cosmology</b> | <b>1</b> |
| <b>3. The stochastic solution</b> | <b>5</b> |
| <b>4. Conclusion</b>              | <b>6</b> |

---

## 1. Introduction

The recent astrophysical observation of the accelerating expansion [1, 2] of the Universe indicates that the cosmological constant  $\Lambda$  has a small nonzero value. However, the origin of the cosmological constant  $\Lambda$  is still unclear [3]. Its origin may be connected, for example, with the presence of some scalar fields [4]. The string inspired gravity is much richer than the Einstein one including several types of scalar fields (dilaton or moduli fields). What is more, the interaction with the background strings may also produce chaos [5]. Similarly, the cosmological constant may gain small stochastic component [6].

The main aim of this work is to examine how small stochastic fluctuation influence the cosmological expansion.

## 2. The classical cosmology

Let us start with the general gravity with some scalar field  $\phi$  and ordinary dustlike matter. The scalar field  $\phi$  may be the remnant of the hybrid inflation model [7] (the “waterfall” field). In the original hybrid inflation model [7] we have two scalar fields: the inflation field  $\varphi$  and the “waterfall” field  $\phi$  described by the potential

$$U(\varphi, \phi) = \frac{1}{2}\lambda(\phi^2 - \phi_0^2) + \frac{1}{2}m^2\varphi^2 + \frac{1}{2}\lambda'\phi^2\varphi^2. \quad (2.1)$$

Fluctuations of the  $\varphi$  late in the inflation period diminishes and  $\phi$  goes to  $\phi_0$ . However, they may be a source of the chaotic fluctuation at the present time. The Lagrange function of the system has correspondingly three terms

$$L = L_g + L_\phi + L_m \quad (2.2)$$

$$L_g = -\frac{1}{2\kappa}R \quad (2.3)$$

$$L_\phi = \frac{1}{2}\partial_\mu\phi\partial^\mu\phi - U(\phi) \quad (2.4)$$

which describes the Einstein gravity  $L_g$ , the “waterfall” field  $L_\phi$  and the dustlike matter  $L_m$ . The scalar potential

$$U(\phi) = U_0(\phi) + \epsilon J_s \phi \quad (2.5)$$

consists of the nonlinear part  $U_0(\phi)$  with a local minimum at  $\phi_0$  and a scalar external source  $J_s$  with a scalar charge  $\epsilon$ . The scalar potential  $U_0(\phi)$  is the remnant from the inflation epoch. The main assumption of this work is that the scalar external source  $J_s$  has the stochastic nature.

The Euler-Lagrange equations of the system (2.2) give the Einstein’s equation

$$R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} = -\kappa T_{\mu\nu} = -\kappa \left( T_{\mu\nu}^m + T_{\mu\nu}^\phi \right) \quad (2.6)$$

where  $\kappa = 8\pi G$  (in  $c = \hbar = 1$  units) together with the equation for the scalar field  $\phi$ . In this equation the fluid is described by the energy-momentum tensor in the standard form  $T_{\mu\nu}^m = (\rho + p)u_\mu u_\nu + Pg_{\mu\nu}$  ( $u^\mu$  is the fluid four-velocity,  $\rho$  is the energy density in the rest frame of the fluid and  $P$  is the pressure in that same frame). For scalar field the energy-momentum tensor has the form  $T_{\mu\nu}^\phi = \partial_\mu \phi \partial_\nu \phi - g_{\mu\nu} L_\phi$ .

The Lagrange-Euler equation of the  $\phi$  field is

$$\square \phi = -\frac{\partial L_\phi}{\partial \phi} = \frac{\partial U_0}{\partial \phi} + \epsilon J_s \quad (2.7)$$

where  $\square f = -\frac{1}{\sqrt{-g}}\partial_\mu(\sqrt{-g}g^{\mu\nu}\partial_\nu f)$  is the d’Alembert operator in a curved space-time. In vacuum (when  $J_s = 0$ ) the static local extremum  $\phi_0$  generates the effective cosmological constant  $\Lambda^0$  with  $\Omega_\Lambda^0 = \rho_\phi^0/\rho_c$ , with  $\rho_\phi^0 = U(\phi_0)$ .

The space-time interval  $ds$  of the Friedman-Robertson-Wolker (FRW) metric is written in the standard way

$$ds^2 = g_{\mu\nu}dx^\mu dx^\nu = c^2 dt^2 - a^2(t) \left[ \frac{dr^2}{1 - kr^2} + r^2 (d\theta^2 + \sin^2 \theta d\phi^2) \right] \quad (2.8)$$

Calculating the Einstein’s equations (2.6) for the metric (2.8) we get two independent equations. The first is known as the Friedman equation,

$$H^2 \equiv \left( \frac{\dot{a}}{a} \right)^2 = \frac{8\pi G}{3} \rho_m + \frac{\Lambda}{3} - \frac{kc^2}{a^2} = \frac{8\pi G}{3} (\rho_m + \rho_\Lambda) - \frac{kc^2}{a^2} \quad (2.9)$$

Here the over-dot denotes a derivative with respect to cosmic time  $t$ . This equation is a constraint equation, in the sense that we are not allowed to freely specify the time derivative  $\dot{a}$ ; it is determined in terms of the energy density and curvature. The second equation, which is an evolution equation, is often combined with the first one (2.9) to obtain the *acceleration equation*

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3} \left( \rho + 3\frac{P}{c^2} \right) + \frac{\Lambda}{3} \quad (2.10)$$

The equation (2.7) for the scalar field  $\phi$  now takes the form

$$\ddot{\phi} + 3\frac{\dot{a}}{a}\dot{\phi} = \frac{\partial U_0}{\partial \phi} + \epsilon J_s \quad (2.11)$$

The substitution

$$a(t) = \exp(\sigma(\tau)) \quad (2.12)$$

transforms the Friedman equation (2.9) into the following equation

$$\left(\frac{d\sigma}{d\tau}\right)^2 = \Omega_m e^{-3\sigma} + \Omega_\Lambda \quad (2.13)$$

where  $t = t_H \tau = H_0^{-1} \tau$  and  $H_0$  is the Hubble constant at the present time  $t_0$  and  $\Omega = \rho/\rho_c$  is the density parameter (in case of the flat universe ( $k = 0$ ) in dust era ( $P = 0$ )). The acceleration equation (2.10) together with the first Friedman equation (2.13) give

$$\frac{d^2\sigma}{d\tau^2} = -\frac{3}{2}\Omega_m e^{-3\sigma} \quad (2.14)$$

Close to the minimum the potential  $U_0(\phi) \sim \frac{1}{2}m_s^2(\phi - \phi_0)^2$  with  $m_s^2 = 2\lambda\phi_0^2$ . Equation for the scalar field (2.7) now takes the form

$$\frac{d^2\phi}{d\tau^2} + 3\frac{d\sigma}{d\tau}\frac{d\phi}{d\tau} = -\frac{1}{H_0^2}\frac{\partial U_0}{\partial\phi} - \epsilon\eta \sim -\mu^2(\phi - \phi_0) - \phi_0\eta \quad (2.15)$$

where

$$\mu = \frac{m_s}{H_0}, \quad \eta = \frac{J_s}{\phi_0 H_0^2} \quad (2.16)$$

is the dimensionless scalar mass and source term. The main assumption of this work is that the scalar field close to the local extremum has partly small stochastic contribution coming from the stochastic source  $\eta$ . In the result the effective cosmological term with a small stochastic contribution originated from the stochastic part of the scalar density  $\delta\rho = \epsilon J_s \phi_0$

$$\Omega_\Lambda = \Omega_\Lambda^0 + \epsilon\lambda\eta \quad (2.17)$$

appears with the dimensionless factor

$$\lambda = \frac{8\pi}{3} \left(\frac{\phi_0}{M_{Pl}}\right)^2 \quad (2.18)$$

In cosmology the equation (2.15) plays the role of the Lagrange equation. The stochastic fluctuations of the cosmological constant will produce the change of scalar field

$$\phi = \phi_0 + \epsilon\phi_0 F(\tau) \quad (2.19)$$

and metric

$$\sigma(t) = \sigma_0(t) + \epsilon \left(\frac{d\sigma_0(\tau)}{d\tau}\right) X(\tau). \quad (2.20)$$

Here  $\epsilon$  is a small perturbation parameter ( $\epsilon \ll 1$ ). The fluctuation of the  $\sigma(t)$  field around the classical  $\sigma_0(t)$  means that also the scale factor will fluctuate around the classical one ( $a_0(t) = \exp(\sigma_0(\tau))$ )

$$a(t) = a_0(t) (1 + \epsilon X(t))$$

together with metric

$$g_{\mu\nu} = g_{\mu\nu}^0 + \epsilon h_{\mu\nu}$$

where

$$h_{\mu\nu} = \begin{pmatrix} 0 & & & \\ -2a_0(t)X(t) & & & \\ & -2a_0(t)X(t) & & \\ & & -2a_0(t)X(t) & \end{pmatrix}$$

describes the space fluctuation around the classical metric

$$g_{\mu\nu}^0 = \begin{pmatrix} 1 & & & \\ & -a_0^2(t) & & \\ & & -a_0^2(t) & \\ & & & -a_0^2(t) \end{pmatrix}$$

The linearization of the equation for the scalar field (2.15) gives

$$\frac{d^2 F}{d\tau^2} + 3 \frac{d\sigma_0(\tau)}{d\tau} \frac{dF}{d\tau} + \mu^2 F = -\eta \quad (2.21)$$

The linearization (eqs. 2.17 and 2.20) using the equation (2.13) leads to the stochastic differential equation

$$\frac{dX}{d\tau} = \frac{\lambda}{2 \left( \frac{d\sigma(\tau)}{d\tau} \right)^2} \eta \quad (2.22)$$

When  $\eta$  represents the white noise ( $\delta\Omega_\Lambda = \eta$ ) its correlation function obeys

$$\langle \eta(\tau)\eta(\tau') \rangle = D\delta(\tau - \tau') \quad (2.23)$$

with  $\delta(\tau)$  being the Dirac distribution.

The solution of the deterministic differential equation

$$\left( \frac{d\sigma_0}{d\tau} \right)^2 = \Omega_m e^{-3\sigma_0(\tau)} + \Omega_\Lambda^0 \quad (2.24)$$

with the present time  $t_0$  ( or  $\tau_0$ ) defined as ( $a_0(\tau_0) = 0$  or  $a_0(t_0) = 1$ )

$$\tau_0 = \frac{t_0}{t_H} = H_0 t_0 = \frac{2}{3\sqrt{\Omega_\Lambda^0}} \text{arcsinh}(\sqrt{\omega_0}) \quad (2.25)$$

and  $\omega_0 = \Omega_\Lambda^0/\Omega_m$  defined by the deterministic cosmological constant  $\Omega_\Lambda^0 = \langle \Omega_\Lambda \rangle$ . The deterministic solution is

$$\sigma_0(\tau) = \frac{2}{3} \ln \left( \frac{1}{\sqrt{\omega_0}} \sinh \left( \frac{3}{2} \sqrt{\Omega_\Lambda^0} \tau \right) \right) \quad (2.26)$$

and

$$a_0(\tau) = \left( \frac{\sinh(\frac{3}{2}\sqrt{\Omega_\Lambda^0}\tau)}{\sqrt{\omega_0}} \right)^{\frac{2}{3}} \quad (2.27)$$

### 3. The stochastic solution

The linear stochastic differential equation (SDE) [8] (2.22) may be rewritten in the following form

$$dX(\tau) = f(\tau)X(\tau)d\tau + d(\tau)dW_\tau \quad (3.1)$$

where  $f(\sigma_0(\tau))$  is the drift coefficient,  $d(\tau)$  is the diffusion coefficient and  $W_\tau$  is a Wiener process which represents intrinsic noise (white noise) ( $\delta\Omega_\Lambda dt \rightarrow dW_t$ ). These differential equations are easy to solve numerically [9]. The formal solution of the equation (3.1) has the following form

$$X(\tau) = \exp\left(\int_{\tau_0}^{\tau} ds f(s)\right) \left(X(\tau_0) + \int_{\tau_0}^{\tau} dW_s \exp\left(-\int_{\tau_0}^s ds f(s)\right)\right) \quad (3.2)$$

The linearized equation (2.22) is the generalized stochastic equation (3.1) with:

$$\begin{aligned} f(\tau) &= 0 \\ d(\tau) &= \frac{\lambda}{2\left(\frac{d\sigma_0(\tau)}{d\tau}\right)^2} \end{aligned} \quad (3.3)$$

where the diffusion coefficient depends straightforward on  $\sigma_0(\tau)$  (2.26). The solution for  $X(\tau)$  takes the simple form:

$$X(\tau) = X(\tau_0) + \frac{\lambda}{2} \int_{\tau_0}^{\tau} \frac{1}{\left(\frac{d\sigma_0(s)}{ds}\right)^2} dW_s$$

The mean value of  $X(\tau)$  takes the form:

$$\langle X(\tau) \rangle = X(\tau_0)$$

According to the equation (2.20) the perturbative solution for  $\sigma(\tau)$  may be rewritten as


$$\begin{aligned} \sigma(\tau) &= \sigma_0(\tau) + \epsilon \frac{d\sigma_0(\tau)}{d\tau} \left( X(\tau_0) + \frac{\lambda}{2} \int_{\tau_0}^{\tau} \frac{1}{\left(\frac{d\sigma_0(s)}{ds}\right)^2} dW_s \right) \\ &= \sigma_0(\tau) + \epsilon \frac{H_0(\tau)}{H_0} \left( X(\tau_0) + \frac{\lambda}{2} \int_{\tau_0}^{\tau} \frac{1}{\left(\frac{d\sigma_0(s)}{ds}\right)^2} dW_s \right) \end{aligned}$$

where  $a_0(\tau) = \exp(\sigma_0(\tau))$  is the deterministic scale factor and  $H_0(\tau) = \dot{a}_0(\tau)/a_0(\tau)$  is the deterministic Hubble parameter.

$$\frac{d\sigma_0(\tau)}{d\tau} = \sqrt{\Omega_\Lambda} \frac{1}{\tanh\left(\frac{3}{2}\sqrt{\Omega_\Lambda}\tau\right)}$$

When the stochastic fluctuations are small enough ( $\delta a(\tau)$ ) we get

$$a(\tau) = a_0(\tau) + \delta a(\tau) = a_0(\tau) + \epsilon a_0(\tau) \frac{H_0(\tau)}{H_0} \left( X(\tau_0) + \frac{\lambda}{2} \int_{\tau_0}^{\tau} \frac{1}{\left(\frac{d\sigma_0(s)}{ds}\right)^2} dW_s \right) + \dots \quad (3.4)$$



**Figure 1:** The scale factor variation with time for deterministic (the black curve) and stochastic expansions (the red one (3.5)). ( $X(\tau_0) = 0$ ,  $\epsilon = 0.5$ ,  $D = 5.0$ ,  $\lambda = 0.5$  for example)

The average calculated with respect to the stochastic process gives

$$\begin{aligned} \langle a(\tau) \rangle = & a_0(\tau) (e^{\epsilon X_0} + \\ & + \frac{1}{4}(\epsilon\lambda)^2 \left( \frac{H_0(\tau)}{H_0} \right)^2 \int_{\tau_0}^{\tau} \frac{ds}{\left( \frac{d\sigma_0(s)}{ds} \right)^2} \int_{\tau_0}^{\tau} \frac{dz}{\left( \frac{d\sigma_0(z)}{dz} \right)^2} \langle \eta(s)\eta(z) \rangle + \dots) \end{aligned}$$

Using (2.23) and assuming that  $X_0 = X(\tau_0) = 0$  we get

$$\langle a(\tau) \rangle = a_0(\tau) \left( 1 + \frac{1}{4}(\epsilon\lambda)^2 D \left( \frac{H_0(\tau)}{H_0} \right)^2 \int_{\tau_0}^{\tau} \frac{ds}{\left( \frac{d\sigma_0(s)}{ds} \right)^4} + \dots \right) \quad (3.5)$$

For late evolutionary times

$$\langle a(\tau) \rangle = a_0(\tau) \left( 1 + \frac{1}{4}(\epsilon\lambda)^2 D \frac{1}{\Omega_{\Lambda}^2} \left( \frac{H_0(\tau)}{H_0} \right)^2 (\tau - \tau_0) + \dots \right)$$

So, if the cosmological constant  $\Lambda$  has a stochastic component  $\delta\Omega_{\Lambda}$  then the average scale factor  $\langle a(\tau) \rangle$  will grow more strongly than in the deterministic case (Fig. 1).

#### 4. Conclusion

In conclusion, the “waterfall” field in the hybrid inflation model as a source of the stochastic process will influence and increase the Universe expansion. It is also a source of the small effective cosmological constant  $\Lambda$  (2.17).

## References

- [1] S.Perlmutter et al. “Supernova Cosmology Project Collaboration”, *Astrophys. J.* **517** (1999) 564, arXiv:astro-ph/9812133, T. Padmanabhan “Advanced Topics in Cosmology: A Pedagogical Introduction”, arXiv:astro-ph/0602117.
- [2] A. G. Riess et all. “Supernova Search Team Collaboration”, *Astron. J.* **116** (1998) 1009, arXiv:astro-ph/9805201.
- [3] For reviews, see S. Weinberg, “The Cosmological Constant Problems”, arXiv:astro-ph/0005265.
- [4] Edmund J. Copeland, M. Sami and Shinji Tsujikawa, “Dynamics of dark energy” arXiv:hep-th/0603057.
- [5] Thibault Damour and Alexander Vilenkin, “Gravitational wave bursts from cosmic strings”, *Phys. Rev. Lett.* **85** (2000) 3761
- [6] J. Martin and M. A. Musso “Stochastic Quintessence”, *Phys. Rev. D* **71** (2005) 063514, arXiv:astro-ph/0410190
- [7] Edmund J Copeland, Andrew R Liddle, David H Lyth, Ewan D Stewart and David Wands, “False Vacuum Inflation with Einstein Gravity”, *Phys. Rev. D* **49** (1994) 6410.
- [8] T.C. Gard, “Introduction to Stochastic Differential Equations”, Marcel Dekker, New York, 1988
- [9] Sasha Cyganowski, Peter E. Kloeden and Jerzy Ombach, “From Elementary Probability to Stochastic Differential Equations with MAPLE”, Springer Verlag (2001).